



General

Guideline Title

ACR Appropriateness Criteria® blunt abdominal trauma.

Bibliographic Source(s)

Sudakoff GS, Rosen MP, Rybicki FJ, Blake MA, Cash BD, Desjardins B, Greene FL, Hindman NM, Oliva IB, Weiss C, Yaghmai V, Expert Panels on Vascular Imaging and Gastrointestinal Imaging. ACR Appropriateness Criteria® blunt abdominal trauma. [online publication]. Reston (VA): American College of Radiology (ACR); 2012. 9 p. [82 references]

Guideline Status

This is the current release of the guideline.

This guideline updates a previous version: Sudakoff GS, Yucel EK, Rosen MP, Francis IR, Baum RA, Foley WD, Gay SB, Greene FL, Mansour MA, Rybicki FJ, Expert Panels on Vascular Imaging, Gastrointestinal Imaging, and Urologic Imaging. ACR Appropriateness Criteria® blunt abdominal trauma. [online publication]. Reston (VA): American College of Radiology (ACR); 2008. 6 p.

Recommendations

Major Recommendations

ACR Appropriateness Criteria®

Clinical Condition: Blunt Abdominal Trauma

Variant 1: Unstable patient.

Radiologic Procedure	Rating	Comments	RRL*
X-ray chest	8	To evaluate for fracture, pneumomediastinum, and abnormal air collection or gas collections, patient condition permitting. Chest radiograph, KUB, and FAST scan are complementary examinations. All are commonly performed in this setting, patient condition permitting.	<input type="text"/>
Single- and two-view plain film (KUB, FAST scan)	1,2,3 Usually appropriate; 4,5,6 May be appropriate; 7,8 Usually not appropriate	To evaluate for free fluid, patient condition permitting. Chest radiograph, KUB, and FAST scan	Relative Radiation

Radiologic Procedure	Rating	Comments	RRL*
		are complementary examinations. All are commonly performed in this setting, patient condition permitting.	
X-ray abdomen and pelvis (KUB)	8	To evaluate for fracture, free intraperitoneal air or abnormal fluid or gas collections. Chest radiograph, KUB, and FAST scan are complementary examinations. All are commonly performed in this setting, patient condition permitting.	<input type="text"/> <input type="text"/>
Arteriography with possible embolization abdomen and pelvis	5	Not appropriate as initial imaging modality but may become more appropriate if additional clinical information or imaging suggests possibility of active hemorrhage.	Varies
CT abdomen and pelvis without contrast	3	Not appropriate for critically unstable patients. Appropriateness rating may increase if clinical condition of patient improves and becomes hemodynamically stable. Would only consider in setting of prior severe contrast reaction or renal failure.	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
CT abdomen and pelvis with contrast	3	Not appropriate for critically unstable patients. Appropriateness rating may increase if clinical condition of patient improves and becomes hemodynamically stable.	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
CT abdomen and pelvis without and with contrast	3	Not appropriate for critically unstable patients.	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
CT chest without contrast	3	Not appropriate for critically unstable patients. Appropriateness rating may increase if clinical condition of patient improves and becomes hemodynamically stable. Would only consider noninfused scanning in setting of prior severe contrast reaction or renal failure.	<input type="text"/> <input type="text"/> <input type="text"/>
CT chest with contrast	3	Not appropriate for critically unstable patients. Appropriateness rating may increase if clinical condition of patient improves and becomes hemodynamically stable.	<input type="text"/> <input type="text"/> <input type="text"/>
CT chest without and with contrast	3	Not appropriate for critically unstable patients. Appropriateness rating may increase if clinical condition of patient improves and becomes hemodynamically stable.	<input type="text"/> <input type="text"/> <input type="text"/>
US abdomen and pelvis	2		O
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Note: Abbreviations used in the tables are listed at the end of the "Major Recommendations" field.

Variant 2: Stable patient.

Radiologic Procedure	Rating	Comments	RRL*
CT abdomen and pelvis with contrast	9	May also consider CT angiography followed by routine portal venous phase sequences if visceral injury (e.g., hepatic, splenic, pancreatic, renal, mesenteric, or vascular injury) suspected clinically or in patients with significant pelvic and/or vertebral fractures.	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
CT chest with contrast	8	May be appropriate for patients who have sustained significant abdominal trauma. The decision of whether to include infused chest CT should be based on the patient's clinical findings and known mechanism of injury.	<input type="text"/> <input type="text"/> <input type="text"/>
CT abdomen and pelvis without and with contrast	7		<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
CT chest without and with contrast	7	The decision of whether to include chest CT should be based on the patient's clinical findings and known mechanism of injury.	<input type="text"/> <input type="text"/> <input type="text"/>
X-ray chest	7		<input type="text"/>
CT chest without contrast	6	The decision of whether to include chest CT should be based on the patient's clinical findings and known mechanism of injury. Consider for patients with known severe contrast allergy or renal failure.	<input type="text"/> <input type="text"/> <input type="text"/>
CT abdomen and pelvis without contrast	6	Consider noninfused CT only if patient has known severe contrast reaction or renal failure (eGFR <40).	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
US chest abdomen and pelvis (FAST scan)	5		O
X-ray abdomen and pelvis (KUB)	5	May be useful to initially evaluate the patient for free intraperitoneal air and/or fractures of the pelvis and vertebral column.	<input type="text"/> <input type="text"/>
Arteriography with possible embolization abdomen and pelvis	4		Varies
US abdomen and pelvis	2		O
<u>Rating Scale:</u> 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Note: Abbreviations used in the tables are listed at the end of the "Major Recommendations" field.

Variant 3: Hematuria >35 RBC/hpf (stable).

Radiologic Procedure	Rating	Comments	RRL*
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate		Recommend a CT program immediately following CT	*Relative Radiation

Radiologic Procedure	Rating	Comments	RRL*
		of abdomen and pelvis in patients with known acute pelvic fractures and/or penetrating injury to the pelvis with hematuria.	<input type="text"/> <input type="text"/> <input type="text"/>
CT chest with contrast	8	The decision of whether to include infused chest CT should be based on the patient's clinical findings and known mechanism of injury.	<input type="text"/> <input type="text"/> <input type="text"/>
X-ray chest	8		<input type="text"/>
CT abdomen and pelvis without and with contrast	7		<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
CT chest without and with contrast	7	The decision of whether to include chest CT should be based on the patient's clinical findings and known mechanism of injury.	<input type="text"/> <input type="text"/> <input type="text"/>
CT chest without contrast	6	The decision of whether to include chest CT should be based on the patient's clinical findings and known mechanism of injury. Consider if patient has known history of severe contrast allergy or renal failure.	<input type="text"/> <input type="text"/> <input type="text"/>
CT abdomen and pelvis without contrast	6	Consider noninfused CT only if patient has known severe contrast reaction or renal failure (eGFR <40).	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
CT pelvis with bladder contrast (CT cystography)	6	Appropriate when performed immediately following CT of abdomen and pelvis. Refer to text for indications.	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
X-ray retrograde urethrography	6	Appropriate in stable patients following blunt or penetrating trauma demonstrating gross blood coming from urethral meatus. Refer to text for indications.	<input type="text"/> <input type="text"/> <input type="text"/>
X-ray abdomen and pelvis (KUB)	5	May be useful to initially evaluate the patient for free intraperitoneal air and/or fractures of the pelvis and vertebral column.	<input type="text"/> <input type="text"/>
X-ray cystography	5	CT cystography preferred to follow initial CT of abdomen and pelvis.	<input type="text"/> <input type="text"/> <input type="text"/>
Arteriography with possible embolization kidney	4		Varies
X-ray intravenous urography	3		<input type="text"/> <input type="text"/> <input type="text"/>
US abdomen and pelvis	3		O
Rating Scale: 1,2,3 Usually not appropriate; 4,5,6 May be appropriate; 7,8,9 Usually appropriate			*Relative Radiation Level

Note: Abbreviations used in the tables are listed at the end of the "Major Recommendations" field.

Summary of Literature Review

Introduction/Background

This review covers only blunt abdominal trauma in adults. Penetrating trauma and pediatric trauma are excluded. In the past 10 years technological advances in multidetector computed tomography (MDCT) have significantly improved the detection of both vascular and visceral injuries following blunt abdominal trauma. With the quicker examination times and improved spatial resolution of current MDCT scanners, sites of parenchymal injury and/or active extravasation are increasingly detected. Detection of extravasation sites from solid organs (e.g., liver, spleen, and kidney), bowel, mesentery, or vascular structures may necessitate therapeutic intervention by either surgery or angiography (with coil embolization or intravascular stent placement). However, not all sites of active extravasation may require intervention, and a trend toward conservative management by trauma surgeons is evolving based not only on the presence of active extravasation but also on its location, its size, and demonstration of enlargement over time. Similar trends are also evolving in the management of isolated peritoneal or pelvic fluid collection following blunt abdominal/pelvic trauma. Small, isolated fluid collections of low Hounsfield density may be observed and managed conservatively, whereas surgical exploration may be considered for larger, higher-density fluid collections.

Hemodynamically Unstable Patients

Hemodynamically unstable patients presenting to the emergency room with clinically obvious major abdominal trauma and with unresponsive profound hypotension need rapid clinical evaluation and immediate resuscitation with volume replacement. If such unstable patients do not respond to resuscitation *and* if they have clear clinical or suspected evidence of abdominal injury, they should go immediately to the operating room without imaging. During resuscitative efforts, if time and circumstances permit, conventional radiographs of the chest and abdomen are often obtained as part of trauma protocols. This may help identify a pneumothorax, pneumoperitoneum, or significant bone injury. Focused abdominal sonography for trauma (FAST) performed by an experienced sonologist to check for intraperitoneal free fluid may quickly provide information that can support a decision to operate immediately, with the caveat that the false negative rate is $\geq 15\%$. More detailed ultrasound (US) to check for organ injury takes too long in this setting and suffers from poor sensitivity.

There is now general agreement that routine diagnostic peritoneal lavage is obsolete because of its invasive nature, lack of specificity, and inability to predict the need for therapeutic surgery. Those patients for whom emergency department resuscitation is successful can be evaluated by contrast-enhanced computed tomography (CT) before surgery. In those patients who cannot be stabilized with fluid and/or pharmacologic intervention, surgery should generally not be delayed by imaging. If the patient becomes hemodynamically stable after surgical intervention, CT scanning with intravenous (IV) contrast should be performed to identify other potential injuries not identified during surgery.

Hemodynamically Stable Patients

Hemodynamically stable patients, patients with mild to moderate responsive hypotension presenting to the emergency room after blunt abdominal trauma, and unstable patients who stabilize after initial resuscitation are in a separate category. They typically have a history of significant trauma and have at least moderate suspicion of intra-abdominal injury based on clinical signs and symptoms. For these patients, two decisions need to be made: 1) Is urgent surgery or therapeutic endoscopic retrograde needed? 2) If not, is a period of close observation warranted? If CT is to be performed, radiographs will offer little if any incremental help with those questions. Rather, the decision to proceed with urgent surgery depends on the identification of specific CT criteria (e.g., active hemorrhage, parenchymal blush or pseudoaneurysm in the spleen, perforation of a hollow viscus, or disruption of the pancreas and/or pancreatic duct, or arterial or venous injury to the aorta or vena cava or their major branches). The decision to operate urgently does not depend solely on the identification of hemoperitoneum or of parenchymal injury to the liver or spleen, because most patients in this category ultimately do not need surgery. However, accurate identification of hemoperitoneum or organ injury is important because patients with these findings require at least a period of close observation. Patients with multiple organ injury or significant active bleeding may require invasive therapy even if they are hemodynamically stable. Conversely, stable patients with isolated organ injury may not need surgery or may need only angiography with embolization even with a large amount of hemoperitoneum.

Ultrasound

Currently US is not an appropriate modality to evaluate these patients, as it may miss up to 25% of liver and spleen injuries, most renal injuries, and virtually all pancreatic, mesenteric, and gut injuries. Multiple studies have shown that US can be insensitive in detecting organ injury, missing up to 62% of spleen and 14% of liver injuries that were found by CT and surgery. US also misses a high proportion of retroperitoneal hemorrhages and of bladder rupture. Combining the results for US in 1,535 abdominal trauma patients from eight published series yields an average US sensitivity of 88% for hemoperitoneum and 74% for organ injury. In a more recent study of 4,029 patients with blunt abdominal trauma, FAST scanning yielded a sensitivity, specificity, and accuracy for the patients with hypotension (87 patients) of 85%, 60%, and 77%, respectively, and in the normotensive group (3,583 patients) yielded a sensitivity, specificity, and accuracy of 85%, 96%, and 95%, respectively. The authors

concluded that hypotensive patients with positive findings with FAST may be triaged directly to surgery and bypass CT depending on the suspected injury. Unfortunately, a negative US (absence of hemoperitoneum) does not rule out significant organ or viscus injury that might require invasive therapy or observation.

US is also insensitive to perforation of gut and to pancreatic injury. For these reasons, it is not very useful in deciding when a patient needs urgent surgery or angiography. For the same reasons, US is not an accurate modality to determine whether a patient needs a period of close observation; thus, if a negative US is the sole imaging modality used to triage a patient, for safety reasons it must be followed by a 12- to 24-hour period of in-hospital observation.

Although US is 63% sensitive to moderate amounts of free intraperitoneal fluid (compared with CT), 400 to 600 cc are needed for US detection of fluid in the trauma setting. Regardless of volume, US diagnosis of free fluid alone does not predict that surgery is needed or that surgery will be therapeutic. In addition, in the best of hands, there is at least a 15% false negative rate for detecting hemoperitoneum with US. US poorly identifies active hemorrhage and also does not accurately predict the need for surgery in splenic injuries. The use of contrast-enhanced US for this purpose is not approved in the United States, but it is increasingly used in monitoring patients suspected of possible solid organ injury following blunt trauma who are considered hemodynamically stable.

Computed Tomography

In hemodynamically stable patients (category B trauma patients), CT accurately predicts whether invasive therapy is urgently needed by identifying active hemorrhage, or hepatobiliary, splenic (either parenchymal contrast blush or pseudoaneurysm), pancreatic, genitourinary, intestinal, or diaphragmatic injury. For these reasons, CT is the primary imaging modality for deciding whether a patient needs urgent surgery, therapeutic angiography, or close observation.

The trend toward placing multidetector CT (MDCT) scanners close to or in emergency departments has substantially diminished the delay in getting patients to the CT scanner and has decreased actual scan time to <40 seconds. In nearly all circumstances, results from MDCT of the abdomen and pelvis can be obtained faster than results from a detailed US of the abdomen or pelvis. In most cases, patient turnaround with rapid-process MDCT can be <10 minutes for a trauma patient.

The radiologist should carefully examine images on the picture archiving and communication system (PACS), or at the CT console, where images can be altered to identify bone injury, pneumoperitoneum, or subtle organ injury. Particular care should be taken to find injury of the spleen because these patients may need observation for potential delayed hemorrhage. In some instances, stable patients with more severe injuries of the liver or spleen plus hemoperitoneum may be managed conservatively with close observation only. It should be noted, however, that various schemes for using CT to grade liver or spleen lacerations are not helpful in deciding whether a patient needs surgery. This decision must be based on the clinical status of the patient in combination with the image findings. If evidence of active hemorrhage is discovered on CT examinations, the patient may undergo arteriography plus embolization or surgery to control the hemorrhage.

The CT image should be carefully examined for subtle signs of pancreatic injury because these patients may need immediate surgery or close observation for signs of complication. Duodenal perforation produces subtle but typical findings on CT, such as extraluminal air or fluid in the retroperitoneum or periportal region. Identifying these findings generally mandates surgical intervention. Duodenal hematoma may not require surgery but does mandate close observation. Other gut injury or perforation produces direct or indirect findings on CT in 50% to 94% of cases. However, if the CT is negative for gut injury in the face of a high clinical suspicion, laparoscopy, surgical exploration, or a period of observation plus repeat CT may be used to further evaluate the patient.

It may also be reasonable to use CT, in conjunction with the clinical information, to decide whether to observe patients in the hospital for a day or send them home after evaluation in the emergency department. The high sensitivity of CT in detecting injuries that require observation in the hospital means that a negative CT may be adequate to release the patient to home in selected cases. US, however, has a substantially lower sensitivity to the kind of injuries that must be observed in the hospital. For this reason, a negative US is not adequate to safely release the patient to home. This weakness of US is reflected in the design of many outcomes-based investigations on the use of US in trauma: all recommend keeping patients with a negative US in the hospital for a period of observation of 12 to 48 hours before release.

There may be a rationale for creating a subcategory of stable patients with trivial trauma, a low clinical index of suspicion, and no signs or symptoms of intra-abdominal injury. In such patients, a negative US alone may be adequate to release them from observation at a lower cost than if CT had been used. CT is necessary, however, if there are any positive findings on US.

Patients with Hematuria

Patients with hematuria after blunt abdominal trauma require some modification to the imaging workup. All patients with gross hematuria and pelvic fracture require additional imaging of the bladder to exclude bladder rupture (absolute indication). A hemodynamically stable patient being

evaluated with MDCT can easily undergo CT cystography using gravity drip infusion of dilute 2% contrast (300 to 500 cc) via an indwelling Foley catheter. Detection of microscopic or gross hematuria without evidence of pelvic fracture or suspected pelvic injury should be considered as a relative, not absolute indication for additional CT cystography.

Identification of clinical indicators of bladder rupture is important when determining if additional bladder imaging is needed in trauma patients with gross hematuria without pelvic fracture or major pelvic injury. Clinical indicators of bladder rupture that may indicate the need for additional bladder imaging with CT or fluoroscopic cystography include suprapubic pain and tenderness, inability to void, low urine output or clots in urine, signs of major perineal trauma such as perineal swelling, hematoma, or blood per meatus. Patients with concomitant head injury, intoxication, altered sensorium, or previous history of bladder outlet obstruction or bladder surgery should be viewed with increased suspicion for bladder injury. If gross blood is identified from the urethral meatus or the prostate is mobile on digital examination (floating), a retrograde urethrogram should be performed first to rule out urethral injury.

CT images should be examined carefully for evidence of renal perfusion, hemorrhage, or extravasation of contrast or urine from the kidney or bladder. All but the worst renal injuries (renal pedicle and pelvis) are generally treated with observation; intraperitoneal bladder rupture is treated with surgical repair. Extraperitoneal bladder rupture is managed with urethral and suprapubic catheter drainage, whereas intraperitoneal bladder rupture requires prompt surgical therapy.

US plays little if any role in the evaluation of genitourinary trauma. Several studies have documented the inability of US to detect injuries of the kidney or bladder in trauma patients.

CT Technique

CT evaluation of the abdomen and pelvis for blunt trauma does not require the use of oral contrast. The use of low or iso-osmolar IV contrast (approximately 110 to 140 cc at 3 to 5 cc per second with a 60 to 70-second scan delay) is essential to identify visceral, vascular, or bowel injury. Scanning includes the lower lung fields through the floor of the pelvis. Delayed imaging through the pelvis (5 minutes) is generally performed if the patient is stable to allow better visualizing of the bladder and distal ureters.

For a CT cystogram, a Foley catheter is placed into the urinary bladder and 500 cc of dilute contrast is instilled into the bladder via the catheter, using the gravity drip technique, after completion of the abdomen and pelvic CT examination. Imaging of the pelvis commences once the rate of bladder contrast has decreased significantly or stopped. At the conclusion of the CT cystogram, the bladder should be drained through the Foley.

Summary

- Hemodynamically unstable patients presenting with blunt abdominal trauma commonly require assessment with chest radiographs, FAST scanning and kidney-ureter-bladder abdominal radiography (KUB).
- MDCT with IV contrast may not be appropriate for patients who are hemodynamically unstable following blunt abdominal trauma.
- MDCT with IV contrast is the imaging modality of choice for evaluating hemodynamically stable patients following blunt abdominal trauma.
- Patients who are hemodynamically stable following blunt (or penetrating) trauma to the abdomen or pelvis with pelvic fracture, gross hematuria, or >35,000 red blood cells per high-power field (RBCs/hpf) require evaluation of the bladder to rule out bladder perforation or urethral injury. This is most easily performed by CT cystography following initial assessment of the abdomen and pelvis with MDCT using IV contrast.

Abbreviations

- CT, computed tomography
- eGFR, estimated glomerular filtration rate
- FAST, focused abdominal sonography for trauma
- hpf, high-power field
- KUB, kidney-ureter-bladder
- RCB, red blood cell
- US, ultrasound

Relative Radiation Level Designations

Relative Radiation Level*	Adult Effective Dose Estimate Range	Pediatric Effective Dose Estimate Range
O	0 mSv	0 mSv
<input type="text"/>	<0.1 mSv	<0.03 mSv

Relative Radiation Level*	Adult Effective Dose Estimate Range 0.1-1 mSv	Pediatric Effective Dose Estimate Range 0.05-0.5 mSv
<div><div></div><div></div><div></div></div>	1-10 mSv	0.3-3 mSv
<div><div></div><div></div><div></div><div></div></div>	10-30 mSv	3-10 mSv
<div><div></div><div></div><div></div><div></div><div></div></div>	30-100 mSv	10-30 mSv
*RRL assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (e.g., region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as “Varies.”		

Clinical Algorithm(s)

Algorithms were not developed from criteria guidelines.

Scope

Disease/Condition(s)

Blunt abdominal trauma

Guideline Category

Diagnosis

Evaluation

Clinical Specialty

Critical Care

Emergency Medicine

Gastroenterology

Internal Medicine

Radiology

Surgery

Urology

Intended Users

Health Plans

Hospitals

Managed Care Organizations

Physicians

Utilization Management

Guideline Objective(s)

To evaluate the appropriateness of initial radiologic examinations for adults with blunt abdominal trauma

Target Population

Adults with blunt abdominal trauma

Note: Penetrating trauma and pediatric trauma are excluded.

Interventions and Practices Considered

1. X-ray
 - Chest
 - Abdomen and pelvis (kidneys, ureters, bladder [KUB])
 - Retrograde urethrography
 - Cystography
 - Intravenous urography
2. Computed tomography (CT)
 - Abdomen and pelvis with contrast
 - Abdomen and pelvis without contrast
 - Abdomen and pelvis without and with contrast
 - Chest without contrast
 - Chest with contrast
 - Chest without and with contrast
 - Pelvis with bladder contrast (CT cystography)
3. Arteriography with possible embolization
 - Abdomen and pelvis
 - Kidney
4. Ultrasound (US)
 - Chest, abdomen, and pelvis (focused abdominal sonography for trauma [FAST] SCAN)
 - Abdomen and pelvis

Major Outcomes Considered

Utility of radiologic examinations in diagnosis of blunt abdominal trauma

Methodology

Methods Used to Collect/Select the Evidence

Searches of Electronic Databases

Description of Methods Used to Collect/Select the Evidence

Literature Search Procedure

The Medline literature search is based on keywords provided by the topic author. The two general classes of keywords are those related to the condition (e.g., ankle pain, fever) and those that describe the diagnostic or therapeutic intervention of interest (e.g., mammography, MRI).

The search terms and parameters are manipulated to produce the most relevant, current evidence to address the American College of Radiology

Appropriateness Criteria (ACR AC) topic being reviewed or developed. Combining the clinical conditions and diagnostic modalities or therapeutic procedures narrows the search to be relevant to the topic. Exploding the term "diagnostic imaging" captures relevant results for diagnostic topics.

The following criteria/limits are used in the searches.

1. Articles that have abstracts available and are concerned with humans.
2. Restrict the search to the year prior to the last topic update or in some cases the author of the topic may specify which year range to use in the search. For new topics, the year range is restricted to the last 5 years unless the topic author provides other instructions.
3. May restrict the search to Adults only or Pediatrics only.
4. Articles consisting of only summaries or case reports are often excluded from final results.

The search strategy may be revised to improve the output as needed.

Number of Source Documents

The total number of source documents identified as the result of the literature search is not known.

Methods Used to Assess the Quality and Strength of the Evidence

Weighting According to a Rating Scheme (Scheme Given)

Rating Scheme for the Strength of the Evidence

Strength of Evidence Key

Category 1 - The conclusions of the study are valid and strongly supported by study design, analysis, and results.

Category 2 - The conclusions of the study are likely valid, but study design does not permit certainty.

Category 3 - The conclusions of the study may be valid, but the evidence supporting the conclusions is inconclusive or equivocal.

Category 4 - The conclusions of the study may not be valid because the evidence may not be reliable given the study design or analysis.

Methods Used to Analyze the Evidence

Systematic Review with Evidence Tables

Description of the Methods Used to Analyze the Evidence

The topic author drafts or revises the narrative text summarizing the evidence found in the literature. American College of Radiology (ACR) staff draft an evidence table based on the analysis of the selected literature. These tables rate the strength of the evidence for all articles included in the narrative text.

The expert panel reviews the narrative text, evidence table, and the supporting literature for each of the topic-variant combinations and assigns an appropriateness rating for each procedure listed in the table. Each individual panel member forms his/her own opinion based on his/her interpretation of the available evidence.

More information about the evidence table development process can be found in the ACR Appropriateness Criteria® Evidence Table Development document (see the "Availability of Companion Documents" field).

Methods Used to Formulate the Recommendations

Expert Consensus (Delphi)

Description of Methods Used to Formulate the Recommendations

Modified Delphi Technique

The appropriateness ratings for each of the procedures included in the Appropriateness Criteria topics are determined using a modified Delphi methodology. A series of surveys are conducted to elicit each panelist's expert interpretation of the evidence, based on the available data, regarding the appropriateness of an imaging or therapeutic procedure for a specific clinical scenario. American College of Radiology (ACR) staff distributes surveys to the panelists along with the evidence table and narrative. Each panelist interprets the available evidence and rates each procedure. The surveys are completed by panelists without consulting other panelists. The ratings are a scale between 1 and 9, which is further divided into three categories: 1, 2, or 3 is defined as "usually not appropriate"; 4, 5, or 6 is defined as "may be appropriate"; and 7, 8, or 9 is defined as "usually appropriate." Each panel member assigns one rating for each procedure per survey round. The surveys are collected and the results are tabulated, de-identified and redistributed after each round. A maximum of three rounds are conducted. The modified Delphi technique enables each panelist to express individual interpretations of the evidence and his or her expert opinion without excessive bias from fellow panelists in a simple, standardized and economical process.

Consensus among the panel members must be achieved to determine the final rating for each procedure. Consensus is defined as eighty percent (80%) agreement within a rating category. The final rating is determined by the median of all the ratings once consensus has been reached. Up to three rating rounds are conducted to achieve consensus.

If consensus is not reached, the panel is convened by conference call. The strengths and weaknesses of each imaging procedure that has not reached consensus are discussed and a final rating is proposed. If the panelists on the call agree, the rating is accepted as the panel's consensus. The document is circulated to all the panelists to make the final determination. If consensus cannot be reached on the call or when the document is circulated, "No consensus" appears in the rating column and the reasons for this decision are added to the comment sections.

Rating Scheme for the Strength of the Recommendations

Not applicable

Cost Analysis

A formal cost analysis was not performed and published cost analyses were not reviewed.

Method of Guideline Validation

Internal Peer Review

Description of Method of Guideline Validation

Criteria developed by the Expert Panels are reviewed by the American College of Radiology (ACR) Committee on Appropriateness Criteria.

Evidence Supporting the Recommendations

Type of Evidence Supporting the Recommendations

The recommendations are based on analysis of the current literature and expert panel consensus.

Benefits/Harms of Implementing the Guideline Recommendations

Potential Benefits

Potential Harms

Focused abdominal sonography for trauma (FAST) performed by an experienced sonologist to check for intraperitoneal free fluid may quickly provide information that can support a decision to operate immediately, with the caveat that the false negative rate is $\geq 15\%$.

Relative Radiation Level (RRL)

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, an RRL indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, both because of organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared to those specified for adults. Additional information regarding radiation dose assessment for imaging examinations can be found in the American College of Radiology (ACR) Appropriateness Criteria® Radiation Dose Assessment Introduction document (see the "Availability of Companion Documents" field).

Qualifying Statements

Qualifying Statements

The American College of Radiology (ACR) Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists, and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the U.S. Food and Drug Administration (FDA) have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Implementation of the Guideline

Description of Implementation Strategy

An implementation strategy was not provided.

Institute of Medicine (IOM) National Healthcare Quality Report Categories

IOM Care Need

Getting Better

IOM Domain

Effectiveness

Timeliness

Identifying Information and Availability

Bibliographic Source(s)

Sudakoff GS, Rosen MP, Rybicki FJ, Blake MA, Cash BD, Desjardins B, Greene FL, Hindman NM, Oliva IB, Weiss C, Yaghmai V, Expert Panels on Vascular Imaging and Gastrointestinal Imaging. ACR Appropriateness Criteria® blunt abdominal trauma. [online publication]. Reston (VA): American College of Radiology (ACR); 2012. 9 p. [82 references]

Adaptation

Not applicable: The guideline was not adapted from another source.

Date Released

1996 (revised 2012)

Guideline Developer(s)

American College of Radiology - Medical Specialty Society

Source(s) of Funding

American College of Radiology (ACR) provided the funding and the resources for these ACR Appropriateness Criteria®.

Guideline Committee

Committee on Appropriateness Criteria, Expert Panels on Vascular Imaging and Gastrointestinal Imaging

Composition of Group That Authored the Guideline

Panel Members: Gary S. Sudakoff, MD (*Principal Author*); Max P. Rosen, MD, MPH (*Panel Chair*, Expert Panel on Gastrointestinal Imaging); Frank J. Rybicki, MD, PhD (*Panel Chair*, Expert Panel on Vascular Imaging); Michael A. Blake, MB, BCh; Brooks D. Cash, MD; Benoit Desjardins, MD, PhD; Frederick L. Greene, MD; Nicole M. Hindman, MD; Isabel B. Oliva, MD; Clifford Weiss, MD; Vahid Yaghmai, MD, MS

Financial Disclosures/Conflicts of Interest

Not stated

Guideline Status

This is the current release of the guideline.

This guideline updates a previous version: Sudakoff GS, Yucel EK, Rosen MP, Francis IR, Baum RA, Foley WD, Gay SB, Greene FL, Mansour MA, Rybicki FJ, Expert Panels on Vascular Imaging, Gastrointestinal Imaging, and Urologic Imaging. ACR Appropriateness Criteria® blunt abdominal trauma. [online publication]. Reston (VA): American College of Radiology (ACR); 2008. 6 p.

Guideline Availability

Electronic copies: Available from the [American College of Radiology \(ACR\) Web site](#) .

Print copies: Available from the American College of Radiology, 1891 Preston White Drive, Reston, VA 20191. Telephone: (703) 648-8900.

Availability of Companion Documents

The following are available:

- ACR Appropriateness Criteria®. Overview. Reston (VA): American College of Radiology; 2 p. Electronic copies: Available in Portable Document Format (PDF) from the [American College of Radiology \(ACR\) Web site](#) .
- ACR Appropriateness Criteria®. Literature search process. Reston (VA): American College of Radiology; 1 p. Electronic copies: Available in PDF from the [ACR Web site](#) .
- ACR Appropriateness Criteria®. Evidence table development – diagnostic studies. Reston (VA): American College of Radiology; 2013 Nov. 3 p. Electronic copies: Available in PDF from the [ACR Web site](#) .
- ACR Appropriateness Criteria®. Radiation dose assessment introduction. Reston (VA): American College of Radiology; 2 p. Electronic copies: Available in PDF from the [ACR Web site](#) .
- ACR Appropriateness Criteria®. Procedure information. Reston (VA): American College of Radiology; 1 p. Electronic copies: Available in PDF from the [ACR Web site](#) .
- ACR Appropriateness Criteria® blunt abdominal trauma. Evidence table. Reston (VA): American College of Radiology; 21 p. Electronic copies: Available in PDF from the [ACR Web site](#) .

Patient Resources

None available

NGC Status

This summary was completed by ECRI on March 19, 2001. The information was verified by the guideline developer on March 29, 2001. This NGC summary was updated by ECRI on January 4, 2006. The updated information was verified by the guideline developer on January 19, 2006. This summary was updated by ECRI Institute on June 2, 2010. This summary was updated by ECRI Institute on September 19, 2012.

Copyright Statement

Instructions for downloading, use, and reproduction of the American College of Radiology (ACR) Appropriateness Criteria® may be found on the [ACR Web site](#) .

Disclaimer

NGC Disclaimer

The National Guideline Clearinghouse® (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion-criteria.aspx>.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.